

Inflation and the dwarf galaxy problem

Jun'ichi YOKOYAMA

*Department of Earth and Space Science, Graduate School of Science,
Osaka University, Toyonaka 560-0043, Japan*

Cosmological N -body simulations with scale-invariant initial perturbations show too many virialized objects of dwarf-galaxy masses in a typical galactic halo in contradiction to observation and stability of galactic discs. We present a model of double hybrid inflation which predicts density fluctuations with a small-scale cutoff to account for the absence of these objects in observations.

PACS Numbers: 98.80.Cq

OU-TAP 142

I. INTRODUCTION

Inflationary cosmology [1,2] has strongly influenced the study of cosmological structure formation for these two decades, because its simplest versions predict a rather specific universe with a vanishingly small spatial curvature and scale-invariant adiabatic fluctuations [3]. The standard models of structure formation today starts with these initial conditions and assume that dominant material contents is cold dark matter (CDM) and a cosmological-constant-like energy of unknown origin.

The standard Λ CDM model has proved remarkably successful in explaining various observational data. In this model the gravitational clustering process is governed by the dark matter component and the baryons do not play important roles. Since gravity is a scale-free force by nature, we would end up with a hierarchical universe if we started with the scale-invariant primordial fluctuations. In such a universe, galaxy halos would appear as scaled versions of clusters of galaxies. That is, just as clusters are associated with thousands of substructure clumps with a steep mass spectrum, we expect many virialized substructures of dwarf-galaxy-scales in galactic halos. Indeed, such substructures are shown to be produced on the relevant scale by recent N -body simulations with higher resolution in the standard Λ CDM model [4,5] in contradiction with observation around our galaxy. Analytic calculations also suggest that galaxies should contain more satellites than observed [6]. One may wonder that this discrepancy is simply due to the fact that these satellites are too dark to be observed. The situation, however, is not that simple because these satellites would affect stability of galactic disks [7]. Hence the problem may not be solved by an adjustment of the mass-to-light ratio.

Recently, Kamionkowski and Liddle [8] proposed that the dearth of substructure in galactic halos might be simply due to the lack of small-scale power of initial fluctuations. They have shown that the discrepancy can be resolved if there is a cutoff at the comoving wavenumber $k \simeq 4.5h\text{Mpc}^{-1} \equiv k_c$ and if the amplitude of smaller-scale perturbation is more than, say, 10^{-2} times smaller than that on larger scale. They have also argued that such a spectrum could be produced if the slope of the inflaton's potential had a discontinuity. While the spectrum of fluctuations produced in such a broken-scale-invariance model has been elegantly studied by Starobinsky [9], there is no particle-physics motivation for considering a scalar potential with its first derivative discontinuous. It would be much better to realize the same feature with a smooth potential.

In the present paper we propose an inflation model with the desired spectrum in which the scalar potential is a smooth function of simple polynomials. We consider a kind of double inflation model [10–12]. Scales probed by large-scale structures and cosmic microwave background (CMB) radiation leave the Hubble radius during the first inflation stage when almost scale-invariant fluctuations with magnitude of order of 10^{-5} are produced, while smaller scales leave the horizon during the second inflation with the amplitude of perturbation smaller than 10^{-7} . Since the spectrum of fluctuations generated near the end of slow-roll inflation models such as new [13] and chaotic [14] inflation deviates from the scale-invariant one severely [11], they cannot be a model for the first inflation. Indeed the desired cutoff scale is so close to those probed by large-scale structure and CMB anisotropy that we must arrange the first inflation to terminate abruptly, in a time scale smaller than the expansion time. We adopt the hybrid inflation model [15] for this purpose. As shown in [15], hybrid inflation can end abruptly with a “waterfall” of a symmetry-breaking scalar field χ , whose potential energy drives inflation while its dynamics is controlled by another scalar field ϕ . We assume that the former field is also coupled to the inflaton field of the second inflation and it sets an appropriate initial condition for the second inflation so that its number of e -folds is controlled in order to ensure the scale of the cutoff lies at the right scale.

The rest of the paper is organized as follows. In §II we describe our model Lagrangian and outline the scenario of cosmic evolution. More detailed analysis and constraints on the model parameters are given in §III. §IV is devoted to discussion and conclusion.

II. MODEL

We consider a double hybrid inflation model. We introduce two complex scalar fields, χ_1 and χ_2 , which induce a series of symmetry-breaking phase transitions, and two real scalar fields, ϕ_1 and ϕ_2 , that control the dynamics of χ 's. The scalar part of our model Lagrangian is given as follows.

$$\mathcal{L} \equiv (\partial\chi_1)^\dagger \partial\chi_1 + \frac{1}{2}(\partial\phi_1)^2 + (\partial\chi_2)^\dagger \partial\chi_2 + \frac{1}{2}(\partial\phi_2)^2 - V(\chi_1, \phi_1, \chi_2, \phi_2), \quad (1)$$

$$V(\chi_1, \phi_1, \chi_2, \phi_2) \equiv V_1(\chi_1, \phi_1) + V_2(\chi_2, \phi_2) + U(\chi_1, \phi_2), \quad (2)$$

$$V_1(\chi_1, \phi_1) = \frac{1}{2}m_1^2\phi_1^2 + \frac{\lambda_1}{2}(|\chi_1|^2 - v_1^2)^2 + g_1^2\phi_1^2|\chi_1|^2, \quad (3)$$

$$V_2(\chi_2, \phi_2) = \frac{1}{2}m_2^2\phi_2^2 + \frac{\lambda_2}{2}(|\chi_2|^2 - v_2^2)^2 + g_2^2\phi_2^2|\chi_2|^2, \quad (4)$$

$$U(\chi_1, \phi_2) = -\frac{1}{2}h^2(|\chi_1|^2 - v_1^2)\phi_2^2 + h^2(|\chi_1|^2 - v_1^2)\sigma\phi_2, \quad (5)$$

where $g_1, g_2, \lambda_1, \lambda_2$ and h are dimensionless coupling constants, and m_1, m_2, v_1, v_2 and σ are constants with mass dimension. Obviously, V_1 and V_2 have the same structure with an appropriate form for hybrid inflation. We assume that V_1 is at higher energy scale than V_2 and U . Then it drives the first inflation and V_2 and U do not affect the dynamics of χ_1 and ϕ_1 in this epoch.

Under the above assumption, evolution of the universe proceeds as follows. The first and the second derivatives of V_1 with respect to χ_1 read

$$\frac{\partial V_1}{\partial \chi_1^\dagger} = \lambda_1(|\chi_1|^2 - v_1^2)\chi_1 + g_1^2\phi_1^2\chi_1, \quad (6)$$

$$\frac{\partial^2 V_1}{\partial \chi_1 \partial \chi_1^\dagger} \equiv M_{\chi_1}^2 = \lambda_1(2|\chi_1|^2 - v_1^2) + g_1^2\phi_1^2, \quad (7)$$

respectively. Hence if $g_1^2\phi_1^2 > \lambda_1 v_1^2$ holds initially, χ_1 has the global minimum at $\chi_1 = 0$. We further take an initial condition such that M_{χ_1} is larger than the Hubble parameter and χ_1 settles to $\chi_1 = 0$ in the expansion time scale. Then the hybrid inflation is driven by the vacuum energy density of χ_1 with some contribution from the mass term of ϕ_1 , which turns out to be small in the late stage of inflation and we neglect it hereafter. Thus the Hubble parameter, H , reads

$$H^2 \cong \frac{8\pi}{3M_{Pl}^2}V_1 \cong \frac{4\pi\lambda_1 v_1^4}{3M_{Pl}^2} \equiv H_1^2. \quad (8)$$

In this regime, the first and the second derivatives of V with respect to ϕ_2 are given by,

$$\frac{\partial V}{\partial \phi_2} = m_2^2\phi_2 + 2g_2|\chi_2|^2\phi_2 + h^2v_1^2(\phi_2 - \sigma), \quad (9)$$

$$\frac{\partial^2 V}{\partial \phi_2^2} \equiv M_{\phi_2}^2 = m_2^2 + 2g_2^2|\chi_2|^2 + h^2v_1^2. \quad (10)$$

Hence it has a minimum at

$$\phi_2 = \phi_{2m} \equiv \frac{h^2(v_1^2 - |\chi_1|^2)}{h^2(v_1^2 - |\chi_1|^2) + 2g_2^2|\chi_2|^2 + m_2^2}\sigma \cong \sigma \quad (11)$$

to which ϕ_2 is anchored provided $M_{\phi_2}^2 > H_1^2$ at $\phi_2 = \phi_{2m}$.

Then, assuming $g_2^2\sigma^2 > \lambda_2 v_2^2$, χ_2 relaxes to $\chi_2 = 0$ if $M_{\chi_2}^2 > H_1^2$. Thus in this period we find $\chi_1 = \chi_2 = 0$ and $\phi_2 = \phi_{2m} \cong \sigma$. On the other hand, ϕ_1 evolves slowly as

$$\phi_1 \propto \exp\left(-\frac{m_1^2}{3H_1^2}H_1 t\right), \quad (12)$$

until it reaches the critical value $\phi_{1c} \equiv \sqrt{\lambda_1}v_1/g_1$ when $\chi_1 = 0$ becomes unstable to trigger the phase transition of χ_1 .

The amplitude of curvature fluctuation generated on comoving scales that leave the horizon during the first inflation is given by

$$\Phi = \frac{3}{5} \frac{H_1}{2\pi\phi_1} \frac{3H_1^2}{m_1^2}, \quad (13)$$

in the matter dominated era, and the spectral index reads

$$n_s = 1 + \frac{2m_1^2}{3H_1^2}. \quad (14)$$

The COBE observation [16], which probes the comoving scale about $e^{5.6}$ times larger than the desired cutoff scale, $2\pi/k_c$, constrains Φ and n_s as

$$\Phi = \frac{9H_1^3}{10\pi\phi_{1*}m_1^2} \simeq 3 \times 10^{-5}, \quad (15)$$

$$n_s = 1.2 \pm 0.3, \quad (16)$$

with

$$\phi_{1*} \equiv \phi_{1c} \exp\left(\frac{5.6m_1^2}{3H_1^2}\right) = \frac{\sqrt{\lambda_1}v_1}{g_1} \exp\left(\frac{5.6m_1^2}{3H_1^2}\right). \quad (17)$$

For $\phi_1 < \phi_{1c}$, the minimum of χ_1 is located on a circle with

$$|\chi_1|^2 = |\chi_{1m}|^2 \equiv v_1^2 - \frac{g_1^2}{\lambda_1} \phi_1^2 = \frac{1}{\lambda_1} M_{\chi_1}^2 (\chi_1 = 0). \quad (18)$$

But χ_1 does not fall to this minimum immediately, because the dynamics of χ_1 is governed by quantum fluctuations intrinsic to an exponentially expanding spacetime during $|M_{\chi_1}^2(\chi_1 = 0)| \lesssim H_1^2$ [17]. Since we need to terminate the first inflation abruptly, we request the time elapsed during $|M_{\chi_1}^2(\chi_1 = 0)| \lesssim H_1^2$ is much smaller than the expansion time H_1^{-1} . Then χ_1 practically traces the evolution of χ_{1m} , where ϕ_1 acquires a large mass,

$$M_{\phi_1}^2 = m_1^2 + \frac{g_1^2}{\lambda_1} M_{\chi_1}^2 (\chi_1 = 0). \quad (19)$$

Thus, unless $g_1^2 \ll \lambda_1$, ϕ_1 rolls down to the origin immediately to terminate the phase transition and χ_1 relaxes to $\chi_1 = v_1$.

In this short period, ϕ_{2m} also shifts from σ to zero. However, it does not change practically until $h^2(v_1^2 - |\chi_{1m}|^2)$ has become smaller than m_2^2 . At this time $M_{\phi_2}^2$ has already decreased to be much smaller than H_1^2 , so that ϕ_2 does not trace the evolution of ϕ_{2m} but remains at σ practically.

Thus after the first inflation both ϕ_1 and χ_1 dissipate their energy through oscillation while χ_2 and ϕ_1 are frozen at 0 and σ , respectively. Then the universe is soon dominated by the vacuum energy density of χ_2 ($= 0$) and the second hybrid inflation sets in. In this regime the amplitude of curvature perturbation must be much smaller than 10^{-5} and the number of e -folds should be around $N_2 \simeq 50 - 55$, so that the cutoff of fluctuation spectrum lies at around $k = 4.5h\text{Mpc}^{-1}$. Here the precise value of N_2 depends on both the scale of the second inflation and on the history of the universe after that, in particular, on whether a mini-inflation like thermal inflation occurs [18]. For definiteness, we take $N_2 = 50$ hereafter.

The second inflation ends as ϕ_2 gets smaller than $\phi_{2c} \equiv \sqrt{\lambda_2}v_2/g_2$ and χ_2 becomes unstable. Hence the maximum amplitude of curvature fluctuations generated in this stage is given by

$$\Phi = \frac{g_2 H_2^3}{\pi \sqrt{\lambda_2} v_2 m_2^2}, \quad (20)$$

measured at the horizon crossing in the radiation era after inflation. It should be smaller than, say, 10^{-7} for our purpose.

The universe is reheated by the decay of χ_2 . Since we do not need to have tiny dimensionless coupling constants in hybrid inflation, the reheat temperature can be much higher than the weak scale to accommodate baryogenesis [19].

After the two inflations and the two phase transitions, we have two networks of cosmic (global) strings, one on comoving horizon scale at the end of inflation with energy scale v_2 and the other on e^{N_2} times larger scale with energy scale v_1 . These strings are cosmologically harmless as long as v_1 and v_2 are smaller than $7 \times 10^{14} \text{GeV}$ [20]. This is why we have assumed χ_1 and χ_2 are complex fields rather than real fields. We would have encountered a fatal domain wall problem in the latter case.

III. DETERMINATION OF MODEL PARAMETERS

Having outlined the evolutionary scenario of the universe in our model, we now consider constraints on the model parameters. First we study constraints obtained from the first and the second inflation in order to produce the desired spectrum in turn. Then we consider the consistency conditions so that evolution of the universe proceeds as outlined in the previous section and that only one field contributes to density fluctuations at each epoch.

A. Constraints on V_1 from the first inflation

Define a dimensionless parameter

$$\alpha_1 \equiv \frac{100m_1^2}{3H_1^2}, \quad (21)$$

then we find

$$\alpha_1 < 10, \quad (22)$$

from (14) and the constraint on the spectral index (16).

Then from the amplitude of fluctuations (15), the scale of the first inflation is determined as

$$v_1 = 4.9 \times 10^{-7} g_1^{-1} \alpha_1 e^{0.056\alpha_1} M_{Pl}, \quad (23)$$

From $v_1 \lesssim 7 \times 10^{14} \text{GeV}$, we find

$$g_1 > 8.4 \times 10^{-3} \alpha_1 e^{0.056\alpha_1}. \quad (24)$$

(21) and (23) determines m_1 as

$$m_1 = 2.7 \times 10^{-13} \lambda_1^{\frac{1}{2}} \alpha_1^{\frac{5}{2}} e^{0.11\alpha_1} M_{Pl}. \quad (25)$$

From the dominance of the vacuum energy density of χ_1 , or

$$\frac{\lambda_1}{4} v_1^4 \gg \frac{1}{2} m_1^2 \phi_{1*}^2, \quad (26)$$

we find

$$g_1 \gg 7.5 \times 10^{-4} \lambda_1^{\frac{1}{4}} \alpha_1^{\frac{3}{4}} e^{0.056\alpha_1}. \quad (27)$$

Since we are assuming that χ_1 has been anchored at the origin by the time ϕ_1 reaches ϕ_{1*} , we require

$$M_{\chi_1}^2 = g_1^2 \phi_{1*}^2 - \lambda_1 v_1^2 + \frac{1}{2} h^2 \sigma^2 > H_1^2 \quad (28)$$

which yields

$$g_1 > 1.0 \times 10^{-6} \alpha_1 e^{0.056\alpha_1} \left(e^{0.056\alpha_1} - 1 \right)^{-\frac{1}{2}}, \quad (29)$$

where contribution of $h^2 \sigma^2 / 2$ is neglected. As ϕ_1 decrease to ϕ_{1c} , $\chi_1 = 0$ becomes unstable and a phase transition is triggered. We require this transition should occur abruptly, that is, the time elapsed from $M_{\chi_1}^2 = H_1^2$ to $M_{\chi_1}^2 = -H_1^2$, which we denote by Δt , should be much smaller than the expansion time. That is,

$$H_1 \Delta t \cong \frac{3H_1^2}{2m_1^2} \ln \left| \frac{\lambda_1 v_1^2 + H_1^2}{\lambda_1 v_1^2 - H_1^2} \right| \cong \frac{3H_1^4}{\lambda_1 v_1^2 m_1^2} = \frac{4\pi}{3} \frac{\lambda_1 v_1^6}{m_1^2 M_{Pl}^4} \ll 1. \quad (30)$$

Here the second approximate equality applies because $v_1 \lesssim 7 \times 10^{14} \text{GeV}$ holds. From (23) and (25), this yields

$$H_1 \Delta t = 1.0 \times 10^{-10} g_1^{-2} \alpha_1 e^{0.11\alpha_1} \ll 1, \quad (31)$$

or

$$g_1 \gg 1.0 \times 10^{-5} \alpha_1^{\frac{1}{2}} e^{0.056\alpha_1}. \quad (32)$$

B. Constraints on V_2 and U from the second inflation

Since ϕ_2 is proportional to $\exp\left(-\frac{m_2^2}{3H_2^2}H_2t\right)$ during the second inflation, we find

$$N_2 = \frac{3H_2^2}{m_2^2} \ln\left(\frac{g_2\sigma}{\sqrt{\lambda_2}v_2}\right) = \frac{2\pi\lambda_2v_2^4}{m_2^2M_{Pl}^2} \ln\left(\frac{g_2\sigma}{\sqrt{\lambda_2}v_2}\right). \quad (33)$$

Let us take the logarithmic factor

$$\ln\left(\frac{g_2\sigma}{\sqrt{\lambda_2}v_2}\right) = 2, \quad (34)$$

for definiteness. Then (20) reads

$$\Phi = 25\sqrt{\frac{2}{27\pi}} \frac{g_2v_2}{M_{Pl}} \equiv 25\sqrt{\frac{2}{27\pi}} \times 10^{-8}\alpha_2 = 3.8 \times 10^{-8}\alpha_2, \quad (35)$$

with α_2 being a constant of order of unity or smaller. Then v_2 is determined as

$$v_2 = 10^{-8}\alpha_2g_2^{-1}M_{Pl}. \quad (36)$$

Now $v_2 \lesssim 7 \times 10^{14}\text{GeV}$ implies

$$g_2 > 1.7 \times 10^{-4}\alpha_2. \quad (37)$$

We also find

$$m_2 = 5.0 \times 10^{-17}\alpha_2^2g_2^{-2}\sqrt{\lambda_2}M_{Pl}, \quad (38)$$

from (33) and

$$\sigma = 1.4 \times 10^{-9}\alpha_2g_2^{-2}\sqrt{\lambda_2}M_{Pl}, \quad (39)$$

from (34). From the dominance of the vacuum energy of χ_2 , $\lambda_2v_2^4/4 \gg m_2^2\phi_2^2/2$, we find

$$g_2 \gg 3.1 \times 10^{-5}\alpha_2^{\frac{1}{2}}, \quad (40)$$

using (36)-(39). Finally, the mass-squared of χ_2 at the origin reads

$$M_{\chi_2}^2(\chi_2 = 0, \phi_2 = \sigma) = (e^4 - 1)\lambda_2v_2^2 = 5.4 \times 10^{-15}\lambda_2\alpha_2^2g_2^{-2}M_{Pl}^2, \quad (41)$$

under our choice of the logarithmic factor (34). This should be larger than the Hubble parameter-squared,

$$H_2^2 = \frac{2\pi\lambda_2v_2^4}{2M_{Pl}^2} = 2.1 \times 10^{-32}\lambda_2\alpha_2^4g_2^{-4}M_{Pl}^2, \quad (42)$$

so that χ_2 is anchored at the origin. This results in a very mild constraint on g_2 ;

$$g_2 > 2.0 \times 10^{-9}\alpha_2. \quad (43)$$

C. Consistency conditions

Here we study constraints on V_2 and U from the first inflation and those on V_1 from the second inflation, so that our scenario works consistently.

During the first inflation each term of V_2 and U should be much smaller than $\lambda_1v_1^4/4$. From $\lambda_1v_1^4/4 \gg \lambda_2v_2^4/4$, we find

$$\Gamma \equiv \frac{\lambda_1v_1^4}{\lambda_2v_2^4} = 5.9 \times 10^6 \frac{\lambda_1g_1^{-4}\alpha_1^4e^{0.22\alpha_1}}{\lambda_2g_2^{-4}\alpha_2^4} \gg 1, \quad (44)$$

and from $\lambda_1v_1^4/4 \gg h^2v_1^2\sigma^2/2$, we obtain

$$\lambda_1g_1^{-2}\alpha_1^2e^{0.11\alpha_1} \gg 8.2 \times 10^{-6}\lambda_2g_2^{-4}\alpha_2^2h^2. \quad (45)$$

We also require $M_{\chi_2}^2 > H_1^2$ and $M_{\phi_2}^2 > H_1^2$ at their respective minima so that these fields are fixed there. The former constraint reads

$$\lambda_2 g_2^{-2} \alpha_2^2 > 4.5 \times 10^{-11} \lambda_1 g_1^{-4} \alpha_1^4 e^{0.22\alpha_1}, \quad (46)$$

while the latter implies

$$h^2 > 1.0 \times 10^{-12} \lambda_1 g_1^{-2} \alpha_1^2 e^{0.11\alpha_1}. \quad (47)$$

On the other hand, χ_1 and ϕ_1 must be fixed at their ground states during the second inflation. Hence we require $M_{\chi_1}^2 (\chi_1 = v_1, \phi_1 = 0) > H_2^2$ and $M_{\phi_1}^2 (\chi_1 = v_1, \phi_1 = 0) > H_2^2$. The former is automatically satisfied because $M_{\chi_1}^2 = \lambda_1 v_1^2 \gg H_1^2 \gg H_2^2$, and the latter turns out to be a very weak constraint on λ_2 and g_2 ;

$$\lambda_2 g_2^{-4} < 2.3 \times 10^{19} \alpha_1^2 \alpha_2^{-4}. \quad (48)$$

D. Determination of model parameters: An example

Having shown all the necessary conditions we now demonstrate there is a wide range of allowed region in the parameter space. So far we have found four lower bounds on g_1 in III A and three lower bounds on g_2 in III B. On the other hand, constraints obtained in III C are more complicated, so let us take $\lambda_1 g_1^{-2} = 1$ and $\lambda_2 g_2^{-2} = 1$ for illustration. These choices are also consistent with our assumption that $g_1^2 (g_2^2)$ is not much smaller than $\lambda_1 (\lambda_2)$.

Then (45) through (48) read, respectively,

$$g_2 \gg 2.9 \times 10^{-3} \alpha_1^{-1} e^{-0.056\alpha_1} \alpha_2 h, \quad (49)$$

$$g_1 \gg 6.7 \times 10^{-6} \alpha_1^2 e^{0.11\alpha_1} \alpha_2^{-1}, \quad (50)$$

$$h > 1.0 \times 10^{-6} \alpha_1 e^{0.056\alpha_1}, \quad (51)$$

$$g_2 > 2.1 \times 10^{-10} \alpha_1^{-1} \alpha_2^2. \quad (52)$$

All of the constraints shown so far are so mild that it is easy to choose a consistent set of parameter values. Here we merely give an example. Let us take Γ not so big, say $\Gamma = 25$, so that the second inflation sets in soon after the end of the first one and that the intermediate period is unimportant. Let us further take $\alpha_1 = 0.1$ and $\alpha_2 = 1$. Then $\Gamma = 25$ implies $g_2 = 0.21 g_1$ under the choice $\lambda_1 g_1^{-2} = \lambda_2 g_2^{-2} = 1$.

Now all the constraints turn out to be those on g_1 and h , which are summarized as

$$g_1 > 8.4 \times 10^{-4}, \quad (53)$$

$$g_1 \gg 1.5 \times 10^{-4}, \quad (54)$$

$$g_1 \gg 0.14h, \quad (55)$$

$$h \gg 1.0 \times 10^{-7}. \quad (56)$$

The scale of inflation then read

$$v_1 = 5.9 \times 10^{11} g_1^{-1} \text{GeV}, \quad (57)$$

$$v_2 = 5.7 \times 10^{11} g_1^{-1} \text{GeV}, \quad (58)$$

together with $\lambda_2 = 4.4 \times 10^{-2} \lambda_1 = 4.4 \times 10^{-2} g_1^2$. Other mass scales of the model are given by

$$m_1 = 10 g_1 \text{TeV}, \quad (59)$$

$$m_2 = 2.9 g_1 \text{TeV}, \quad (60)$$

$$\sigma = 8.0 \times 10^{10} g_1^{-1} \text{GeV}. \quad (61)$$

If we take $g_1 = 0.1$, for example, all the dimensionless coupling constants take natural values and v_1 and v_2 fall on the typical symmetry breaking scale for the axion [21].

IV. CONCLUSION

In the present paper we have presented a simple model of double inflation that can realize a spectrum of density fluctuation with a small-scale cutoff at the comoving wavenumber at $k_c \simeq 4.5 h \text{Mpc}^{-1}$ in order to explain the dearth of substructure in galactic halos. Since the spectrum of density fluctuations generated in the late stage of new and chaotic inflation models deviates from the scale-invariant one significantly, we adopted the hybrid inflation model for the first inflation which is responsible for fluctuations on scales probed by large-scale structures and CMB anisotropy. We have also made use of the hybrid inflation model for the second inflation because it is easy to control the energy scale of inflation in this model.

Since each hybrid inflation requires at least two scalar fields, our model includes four scalar fields as a double hybrid inflation model. In this sense our model is somewhat complicated, but our scalar potential consists of a polynomial up to only fourth order and all the parameters take natural magnitudes. There are, however, two unwanted features in this model. In concluding the paper, we point them out and mention possible resolutions in turn.

One is the problem of initial condition. It has been discussed that only a tiny fraction in the field configuration space can give rise to successful hybrid inflation, because ϕ rolls down the potential rapidly if χ is too large initially [22]. The resolution to this problem has recently been proposed [23], which introduces more fields or brane cosmology.

The other is the linear interaction term of ϕ_2 in (5), which is required in order to shift the potential minimum of ϕ_2 out of $\phi_2 = 0$ during the first inflation. There is no particle-physics motivation to introduce this type of direct linear coupling. But such a term is effectively induced naturally in more complicated classes of supergravity inflation models [12].

The purpose of the present paper was to make the simplest possible toy model to realize the desired spectral shape of primordial density fluctuations in order to stress the essential ingredients of the model. In a forthcoming paper we present a more realistic supergravity model to realize the same spectral feature.

ACKNOWLEDGMENTS

The author is grateful to F. Takahara for useful communications. This work was partially supported by the Monbusho Grant-in-Aid, Priority Area “Supersymmetry and Unified Theory of Elementary Particles” (#707) and the Monbusho Grant-in-Aid for Scientific Research No. 11740146.

-
- [1] A.H. Guth, Phys. Rev. **D23**, 347 (1981); K. Sato, Mon. Not. R. astr. Soc. **195**, 467 (1981).
 - [2] For a review of inflation see, *e.g.* A.D. Linde, Particle Physics and Inflationary Cosmology (Harwood, 1990); D.H. Lyth and A. Riotto, Phys. Rep. **314**, 1(1999).
 - [3] S.W. Hawking, Phys. Lett. **115B**, 295 (1982); A.A. Starobinsky, Phys. Lett. **117B**, 175 (1982); A.H. Guth and S-Y. Pi, Phys. Rev. Lett. **49**, 1110 (1982).
 - [4] A.A. Klypin, A.V. Kravtsov, and O. Valenzuela, Astrophys. J. **522**, 82 (1999).
 - [5] B. Moore, et al. Astrophys. J. **524**, L19 (1999).
 - [6] G. Kauffmann, S.D.M. White, and B. Guiderdoni, Mon. Not. R. astr. Soc. **264**, 201 (1993).
 - [7] G. Toth and J.P. Ostriker, Astrophys. J. **389**, 5 (1992).
 - [8] M. Kamionkowski and A.R. Liddle, Phys. Rev. Lett. **84**, 4525 (2000).
 - [9] A.A. Starobinsky, JETP Lett. **55**, 489(1992).
 - [10] J. Silk and M.S. Turner, Phys. Rev. **D35**, 419 (1987); L.F. Kofman, A.D. Linde, and A.A. Starobinsky, Phys. Lett. B **157**, 361 (1985); R. Holman, E.W. Kolb, S.L. Vadas, and Y. Wang, Phys. Lett. B **269**, 252 (1991); D. Polarski and A.A. Starobinsky, Nucl. Phys. **B385**, 623 (1992); Phys. Rev. **D51**, 6123 (1994); S. Gottlöber, J.P. Muecket, and A.A. Starobinsky, Astrophys. J. **434**, 417 (1994); L. Randall, M. Soljacić, and A.H. Guth, Nucl. Phys. **B472**, 377 (1996); J. García-Bellido, A.D. Linde, and D. Wands, Phys. Rev. **D54**, 6040 (1996); J. Yokoyama, Phys. Rev. **D58**, 083510 (1998).
 - [11] J. Yokoyama, Phys. Rev. **D59**, 107303 (1999).
 - [12] K. Izawa, M. Kawasaki, and T. Yanagida, Phys. Lett. B **411**, 249 (1997); M. Kawasaki, N. Sugiyama, and T. Yanagida, Phys. Rev. **D57**, 6050 (1998); M. Kawasaki and T. Yanagida, Phys. Rev. **D59**, 043512 (1999); T. Kanazawa, M. Kawasaki, N. Sugiyama, and T. Yanagida, Phys. Rev. **D61**, 023517 (1999); T. Kanazawa, M. Kawasaki, and T. Yanagida, Phys. Lett. B **482**, 174 (2000); T. Kanazawa, M. Kawasaki, N. Sugiyama and T. Yanagida, astro-ph/0006445.
 - [13] A.D. Linde, Phys. Lett. **108B**, 389 (1982); A. Albrecht and P.J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).
 - [14] A.D. Linde, Phys. Lett. **129B**, 177 (1983).
 - [15] A.D. Linde, Phys. Lett. B **259**, 38 (1991); Phys. Rev. **D49**, 748 (1994).
 - [16] C.L. Bennett et al. Astrophys. J. **464**, L1(1996).

- [17] E.T. Vishniac, K.A. Olive, and D. Seckel, Nucl. Phys. **B289**, 717 (1987); J. Yokoyama, Phys. Rev. Lett. **63**, 712 (1989); M. Nagasawa and J. Yokoyama, Nucl. Phys. B **370**, 472 (1992).
- [18] D. H. Lyth and E. D. Stewart, Phys. Rev. Lett. **75**, 201 (1995); D. H. Lyth and E. D. Stewart, Phys. Rev. **D53**, 1784 (1996).
- [19] For a review of non-GUT baryogenesis, see *e.g.* A.D. Dolgov, Phys. Rept. **222**, 311 (1992).
- [20] A. Vilenkin and E.P.S. Shellard, Cosmic Strings and Other Topological Defects (Cambridge, 1994).
- [21] For the latest constraint on the symmetry breaking scale of the axion, see M. Yamaguchi, M. Kawasaki, and J. Yokoyama, Phys. Rev. Lett. **82**, 4578 (1999).
- [22] N. Tetradis, Phys. Rev. **D57**, 5997 (1998).
- [23] L.E. Mendes and A.R. Liddle, astro-ph/0006020.